

Table 1

Metribuzin Usage

Site	Acres (000) Planted	Acres Treated (000)		Lb AI Applied (000)		Application Rates			States of Most Usage and % of Usage in These States
		-----		-----		-----			
		Likely Average	Likely Max	Likely Average	Likely Max	lb ai/# acre/yr	appl lb /year	ai/ A app 1	
Corn	76,200	180	290	30	80	0.2	1.0	0.2	IA 99%
Barley	8,200	5	10	3	15	0.6	1.0	0.6	WA UT 85%
Wheat	71,500	400	500	65	80	0.2	1.0	0.2	WA OR 92%
Sorghum	11,600	1	4	1	3	0.6	1.0	0.6	IA 90%
Lentils	130	<1	<1						
Peas, Dry	170	60	100	15	25	0.3	1.0	0.3	ID WA 90%
Peas, Green	320	15	30	4	7	0.2	1.0	0.2	WA OR 92%
Alfalfa	24,800	200	340	110	200	0.6	1.0	0.6	WA OR UT ID MT MI 86%
Hay, other	36,000	10	30	4	12	0.4	1.0	0.4	NJ WA 85%
Potatoes	1,400	830	920	430	550	0.5	1.0	0.5	ID WA WI ME FL OR 77%
Soybeans	59,300	6,540	11,540	1,980	3,690	0.3	1.0	0.3	OH IL IN IA MO MI 66%
Sugarcane	900	90	140	90	180	1.0	1.0	1.0	LA FL 100%
Grasses & Turf		30	60	30	60	1.0	1.0	1.0	
Asparagus	90	25	50	25	51	1.1	1.0	1.1	WA CA 81%
Carrots	100	5	10	1	2	0.2	1.0	0.2	MN WI 81%
Tomatoes	450	110	130	60	80	0.5	1.0	0.5	FL OH MI IN TN PA 78%
Sainfoin	NO DATA								

NOTES
Calculation:; of the **above** numbers may not **appear** to agree because they are displayed as rounded
% of each crop treated with metribuzin is 1%or less, except for soybeans. The likely average % of soybeans treated
is 11% and the likely maximum % of soybeans treated is 20%.